



HINCHINBROOK 2011-2020 2020 REVIEW

Tagged fish numbers, catch rates, fish sizes, growth, locations and recaptured fish

The following dashboard provides a summary of data for the Hinchinbrook area from Cardwell to Ingham covering the past 10 years.

Summary

- From 2011-2020 there were 6,074 fish tagged for 207 recaptures (3.4%) in the Hinchinbrook area covered by grid maps HCC, HIC and IGH
- Key species tagged were Barramundi 23.5%, Blackspotted Rockcod 13.9% and Mangrove Jack 13.7%, Goldspotted Rockcod 13.4% and Pikey Bream 13.4% accounting for 78.0% of all fish tagged
- The catch rate was calculated as the number of fish tagged/fisher day and the rate in 2020 was 3.8 which is a little above the 10 year average historic catch rate of 3.6 fish/fisher day
- Of the 203 recaptures with the postcode of the fisher recorded 87 (42.7%) were local from Ingham-Cardwell
- The remaining 116 recapture fishers were from the Cairns area in the north to South East Qld in the south with 16 (7.9%) being interstate fishers mostly from NSW
- Postcodes of recaptures suggest that the Hinchinbrook area is attractive for visiting fishers
- The average length of Barramundi in 2020 was 490mm and ranged from 330-820mm with the average length over the past 10 years being below legal size of 580mm
- No trophy fish 1+m were tagged over the past 10 years
- The percentage of legal Barramundi in 2020 was 21.9%
- Over the past 10 years 42.7% of Barramundi recaptures were in the same area as tagged with 6 fish being recaptured outside the Hinchinbrook area
- The average length of Mangrove Jack in 2020 was 346mm and has been close to the minimum legal length of 350mm for the past 10 years
- Of the 19 Mangrove Jack recaptures 14 (73.7%) were recaptured in the same area as tagged with 1 fish recaptured offshore 50km east
- Blackspotted and Goldspotted Rockcod data were combined as they are closely related species with similar habits
- The average length of Rockcod in 2020 was 327mm and had been below the minimum legal length of 380mm for the past 10 years with just 13 trophy fish over 600mm in that time
- Of the 92 Rockcod recaptures 79 (85.9%) were recaptured in the same area as tagged with the longest distance moved being 22km and no recaptures recorded outside the area

Infofish Australia January 2021



HINCHINBROOK	BARRAMUNDI MANGROVE JACK	GOLDSPOTTED ROCKCOD BLACKSPOTTED ROCKCOD
Induces Ind		
MEASURE	ACTUAL	GRAPH or MAP
Cardwell to Ingham Hinchinbrook Channel Hinchinbrook Island Seymour River Victoria Creek	Grid maps HCC HIC IGH	nen konste Tursen ander
Catch and effort number of tagged fish fisher tag days Taggers 2011-2020 Tag rate	HIGHEST LOWEST NOW TAGGED FISH 2017 2020 979 319 458 EFFORT 2011 2017 2020 2011 2017 2020 119 HIGHEST LOWEST NOW 100	HINCHINBROOK TAG AND EFFORT 1200 1000 1
Fish/fisher day 2011-2020	TAG RATE 2015 2017 2020 4.6 3.0 3.8 10 YEAR AVERAGE 3.6	50 45 40 42 30 42 42 42 42 42 42 42 42 42 42 42 42 42
Fish tagged and recaptured 2011-2020	TAGGEDRECAPTURED6074207RECAPTURE RATE3.4%	FISH TAGGED AND RECAPTURED HINCHINBROOK
Species composition Numbers tagged 2011-2020	BARRAMUNDI 1,427 BLACKSPOTTED/GOLDSPOTT ED ROCKCOD 846/816 MANGROVE JACK 835 PIKEY BREAM 812	HINCHINBROOK SPECIES COMPOSITION

Species composition	BARRAMUNDI	HINCHINBROOK SPECIES PERCENTAGE
Percentage tagged	23.5%	
2011-2020	BLACKSPOTTED/GOLDSPOTT	60%
	ED ROCKCOD	
	13.9%/13.4%	
	MANGROVE JACK	2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 BARRAMUNDI BS ROCKCOD MANGROVE JACK GS ROCKCOD
	13.7%	PIKEY BREAM BARRED JAVELIN GOLDEN SNAPPER OTHER
	PIKEY BREAM	
	13.4%	
Postcodes of recapture	POSTCODES FISHERS	RECAPTURE FISHER POSTCODES
fishers (203 with	INGHAM-CARDWELL 87	CENTRAL QLD
postcodes)	TOWNSVILLE 37	OTHER STATES
2011-2020	SE QLD 32	SOUTH EAST QLD
	TULLY-CAIRNS 20	TOWNSVILLE TO MACKAY
	OTHER STATES 16	0 20 40 60 80 100
	CENTRAL QLD 11	RECAPTURES

MEASURE	ACTUAL	GRAPH or MAP
Barramundi		
Average length (mm) Bars show smallest to largest fish 2011-2020	HIGHEST LOWEST NOW AVERAGE LENGTH 2020 2015 2018 2020 570 518 490 LARGEST 2015 2014 2020 970 670 820	HINCHINBROOK BARRAMUNDI AVERAGE LENGTH
Fish lengths percentage Undersize <580mm Legal 580-1000mm Trophy 1000+mm 2011-2020	970 670 820 HIGHEST LOWEST NOW LEGAL 2017 2014 2020 41.4 9.3 21.9 TROPHY 2011-2019 2020 0 0 0	HINCHINBROOK BARRAMUNDI LENGTHS
Barramundi growth at the end of each year based on all recaptures since 2011 (30)	GROWTHYEAR 195mmYEAR 2158mmYEAR 3195mmYEAR 4221mmYEAR 5242mm	HINCHINBROOK BARRAMUNDI GROWTH
Grid locations of tagged fish (maps HCC, HIC, IGH) 2011-2020	Grids where most fish caught GRID FISH	
Barramundi movements Hinchinbrook area (45) Distance moved (km) 2011-2020	Recaptured same area 21 (46.7%) Longest distance north 53 Longest distance south 175	HINCHINBROOK BARRAMUNDI MOVEMENT
Barramundi recaptured outside Hinchinbrook area (6) Recapture location 2011-2020	North to Tully River 4 South to Townsville 2	

MEASURE	ACTUAL	GRAPH or MAP
Mangrove Jack		
Average length (mm) Bars show smallest to largest fish 2011-2020	HIGHEST LOWEST NOW AVERAGE LENGTH 2020 366 316 346 LARGEST 2015 2016 2020 500 445 440	HINCHINBROOK MANGROVE JACK AVERAGE LENGTH TROPHY 4000 4000 4000 4000 4000 4000 4000 40
Fish lengths percentage Undersize <350mm Legal 350-500mm Trophy 500+mm 2011-2020	HIGHEST LOWEST NOW LEGAL 2016 2012 2020 67.1 37.1 54.7 TROPHY 2015 2016-19 2020 1.0 0 0	HINCHINBROOK MANGROVE JACK LENGTHS
Mangrove Jack growth at the end of each year based on all recaptures since 2011 (14)	Insufficient recaptures	HINCHINBROOK MANGROVE JACK GROWTH
Grid locations of tagged fish (map HCC, HIC, IGH) 2011-2020	Grids where most fish caught GRID FISH	
Mangrove Jack movements Hinchinbrook area Distance moved (km) 2011-2020	Recaptures 19 Recaptured same area 14 (73.7%) Longest distance moved 50	HINCHINBROOK MANGROVE JACK MOVEMENT
Mangrove Jack recaptured outside Hinchinbrook area 2011-2020	Recaptures 1 Moved 50km offshore	

MEASURE	ACTUAL	GRAPH or MAP
Blackspotted Rockcod Goldspotted Rockcod		
Average length (mm) Both species Bars show smallest to largest fish 2011-2020 Fish lengths percentage Both species	HIGHEST LOWEST NOW AVERAGE LENGTH 2020 348 298 327 LARGEST 2015 2020 1050 470 470 HIGHEST LOWEST NOW LEGAL LOWEST NOW	HINCHINBROOK ROCKCOD AVERAGE LENGTH
Undersize <380mm Legal 380-600mm Trophy 600+mm 2011-2020 Both species growth at the end of each year	2011 2013 2020 31.2 9.4 25.0 TROPHY 2016 2019 2020 3.5 0 0 Blackspotted Rockcod YEAR 1 77mm	806 705 507 507 507 507 507 507 507
based on all recaptures Blackspotted Rockcod (43) Goldspotted Rockcod (17) since 2011	YEAR 2 126mm YEAR 3 156mm YEAR 4 176mm YEAR 5 192mm Goldspotted Rockcod Insufficient data	200 100 100 100 0 0 0 0 0 0 0 0 0 0 0 0
Grid locations of tagged fish (maps HCC, HIC, IGH) 2011-2020	Grids where most fish caughtGRIDFISHIGH-U52184HCC-V63141HCC-06590HCC-P6483	Carrier Carrie
Rockcod movements Hinchinbrook area Blackspotted Rockcod (53) Goldspotted Rockcod (38) Distance moved (km) 2011-2020	Recaptured same area 79 (85.9%) Longest distance moved 22 No recaptures outside Hinchinbrook area	HINCHINBROOK ROCKCOD MOVEMENT

MEASURE	ACTUAL	GRAPH or MAP
Pikey Bream		
Average length (mm) Bars show smallest to largest fish 2011-2020	HIGHESTLOWESTNOWAVERAGELENGTH2020201820112020259236253LARGEST20132019201320192020400300350	HINCHINBROOK PIKEY BREAM AVERAGE LENGTH 450.0 100.0 100.0 100.0 100.0 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020
Fish lengths percentage Undersize <250mm Legal 250-399mm Trophy 400+mm 2011-2020 Growth at the end of each year based on all	400 300 350 HIGHEST LOWEST NOW LEGAL 2014 2020 75.8 38.7 37.0 TROPHY 2013 2014-19 2013 2014-19 2020 1.1 0 0 Insufficient data Recaptures	HINCHINBROOK PIKEY BREAM LENGTHS
recaptures since 2011 Grid locations of tagged fish (maps HCC, HIC, IGH) 2011-2020	Grids where most fish caughtGRIDFISHIGH-Y5354IGH-AD4642HCC-P6440	sie Cett
Pikey Bream movements Hinchinbrook area Distance moved (km) 2011-2020	Recaptures 17 Recaptured same area 11 (64.7%) Longest distance moved 28 No recaptures outside Hinchinbrook area	
Product of Infofis	CRYSTALBOWL	Supported by Suntag

Data sources

Data used in the Hinchinbrook Crystal Bowl are:

• Tag records and recapture records at http://crystalbowl.infofishaustralia.com.au/

Basis for analysis

- Catch rates calculated as number of fish tagged/fisher/day for taggers with a minimum of 12 fisher days (at least 1 per month) to provide a reliable catch rate
- For Barramundi undersized fish are <580mm, legal fish are >=580 and <1000mm and trophy fish are >=1000mm
- For Mangrove Jack undersized fish are <350mm, legal fish are >=350 and <500mm and trophy fish are >=500mm
- For Rockcod undersized fish are <380mm, legal fish are >=380 and <600mm and trophy fish are >=600mm
- Analysis of Goldspotted Rockcod and Blackspotted Rockcod is based on combined records as the species are closely related
- Growth calculated using logarithmic curve from recaptures where fish were out for 90 days or more and had positive growth
- Locations based on Infofish grid maps HCC, HIC and IGH and other grid maps where fish were recaptured outside the area
- Movement has been calculated as the shortest distance by water from the tag to the recapture location derived from Google Earth